

University of Groningen

Copper-catalyzed enantioselective synthesis of trans-1-Alkyl-2-substituted cyclopropanes via tandem conjugate addition-intramolecular enolate trapping

Den Hartog, T.; Rudolph, A.; Macia, B.; Minnaard, A.J.; Feringa, B.L.

Published in:
Journal of the American Chemical Society

DOI:
[10.1021/ja105704m](https://doi.org/10.1021/ja105704m)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2010

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Den Hartog, T., Rudolph, A., Macia, B., Minnaard, A. J., & Feringa, B. L. (2010). Copper-catalyzed enantioselective synthesis of trans-1-Alkyl-2-substituted cyclopropanes via tandem conjugate addition-intramolecular enolate trapping. *Journal of the American Chemical Society*, 132(41), 14349 - 14351. <https://doi.org/10.1021/ja105704m>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Copper-Catalyzed Enantioselective Synthesis of *trans*-1-Alkyl-2-substituted Cyclopropanes via Tandem Conjugate Addition–Intramolecular Enolate Trapping

Tim den Hartog, Alena Rudolph, Beatriz Maciá, Adriaan J. Minnaard,* and Ben L. Feringa*

Stratingh Institute for Chemistry, University of Groningen, Nijenborgh 4, 9747 AG Groningen, The Netherlands


Received June 29, 2010; E-mail: b.l.feringa@rug.nl; a.j.minnaard@rug.nl

Abstract: Cu-TolBINAP-catalyzed conjugate addition of Grignard reagents to 4-chloro- α,β -unsaturated esters, thioesters, and ketones leads to 4-chloro-3-alkyl-substituted thioesters and ketones in up to 84% yield and up to 96% ee upon protonation of the corresponding enolates at low temperature. Tandem conjugate addition–enolate trapping, however, yields *trans*-1-alkyl-2-substituted cyclopropanes in up to 92% yield and up to 98% ee. The versatility of this reaction is illustrated by the formation of key intermediates for the formal syntheses of cascarillic acid and grenadamide.

The enantioselective Simmons–Smith cyclopropanation with a stoichiometric amount of a chiral dioxaborolane, introduced by Charette and co-workers,¹ is currently the benchmark reaction for the synthesis of *trans*-1-alkyl-2-substituted cyclopropanes. Only a few catalytic² asymmetric methods^{3,4} can compete with this method¹ in terms of efficiency and yield. This is in sharp contrast to the well-known catalytic asymmetric synthesis of *trans*-1-aryl-2-substituted cyclopropanes.^{5,6} Although Michael addition-initiated ring-closure reactions (MIRCs) are well-known for the diastereo- and enantioselective preparation of substituted cyclopropanes,⁵ to the best of our knowledge the use of organometallic reagents for the catalytic asymmetric^{6,7} preparation of *trans*-1,2-disubstituted cyclopropanes^{6a,b,e,1} via MIRC is unprecedented in the literature.⁸

Recently, we explored the asymmetric allylic alkylation⁹ of 4-halocrotonates using Grignard reagents for the preparation of α -Me-substituted ester **2** in high yield and high regio- and enantioselectivity (Scheme 1a, R¹ = OBn).¹⁰ Here we report the Cu-catalyzed asymmetric 1,4-addition (ACA) of Grignard reagents^{9b,c,11} to 4-chlorocrotonates using the readily available TolBINAP as ligand. This transformation provides a route to 4-chloro-3-alkyl-substituted thioesters and ketones (Scheme 1b, **4**, X = Cl) and a general method to form *trans*-1,2-disubstituted cyclopropanes (Scheme 1c, **5**, X = Cl) with excellent enantioselectivities.

Scheme 1. Switch in Regioselectivity for the Reactions of 4-Halocrotonates with Grignard Reagents



To synthesize *trans*-cyclopropanes via tandem ACA–enolate trapping, initially the ACA of Grignard reagents to 4-halocrotonates was studied while varying the ligand (Table 1). The low yield observed¹² for the ACA to bromo-substituted thioester **6a** (entry 1) prompted us to investigate the chloro-substituted analogue **6b** for this reaction (entry 2). Addition of hexylmagnesium bromide to **6b** using Cu-TolBINAP at –78 °C gave, after quenching at this temperature, 4-chloro-3-hexyl thioester **7b** in good yield (83%) and excellent ee (94%). For 4-halo- α,β -unsaturated aliphatic ketones and esters, similar results were obtained. For instance, reactions of the bromo-substituted substrates **6c** and **6d** with Grignard reagents gave low yields of, respectively, cyclopropanes **8c** and **8d** (entries 3 and 4). In contrast, the reaction of the 4-chloro-substituted α,β -unsaturated ketone **6e** gave **7e** in good yield (entry 5, 84%) and excellent ee (96%). The Cu-catalyzed reaction of ester **6f** with phenethylmagnesium bromide at –40 °C gave cyclopropane **8f** as the main product, along with traces of **7f**, in a reasonable yield (65%, entry 6) and excellent ee (>95%). Already at –40 °C, the enolate formed from **6f** undergoes slow ring-closure to yield the corresponding cyclopropane, **8f**.


Table 1. Substrate Screening for ACA to 4-Halocrotonates^a

Reaction scheme showing the asymmetric allylation of ketones using R^2MgBr , CuI , and (R) -ToIBINAP in $tBuOMe/CH_2Cl_2$ at $-78\text{ }^\circ C$ for 16 h. The starting material **6** (an allyl ketone with leaving group LG) reacts to form product **7** (a linear ketone) or product **8** (a cyclic ketone). The chiral ligand **L2** ((R) -ToIBINAP) is shown as a binaphthyl derivative with two $PpTol_2$ groups.

entry	LG		R ¹	R ²	product	yield (%)	ee ^b (%)
1	Br	6a	SEt	hexyl	8a	<20 ^c	—
2	Cl	6b	SEt	hexyl	7b	83	94
3	Br	6c	C ₁₁ H ₂₃	hexyl	8c	32 ^d	n.d.
4 ^e	Br	6d	OMe	BnCH ₂	8d	<20 ^c	—
5	Cl	6e	C ₁₁ H ₂₃	BnCH ₂	7e	84	96
6 ^e	Cl	6f	OMe	BnCH ₂	8f	65 ^g	>95

^a Conditions: **6** (0.5 mmol) in CH₂Cl₂ added over 2 h to a solution of CuI (1 mol %), TolBINAP (1.5 mol %), and Grignard reagent (1.2 equiv) in *t*BuOMe. ^b Determined by chiral GC or HPLC. ^c Based on GC–MS. ^d 20% of **6c** and 18% of (*E*)-pentadec-2-en-4-one obtained. ^e Reaction performed at –40 °C, 4 h reaction time. ^f In addition, traces (<10%) of **7f** were obtained; in all other entries, exclusively **7** or **8** was obtained. ^g Combined yield of **7f** and **8f**.

The formation of cyclopropanes via tandem ACA–enolate trapping was subsequently investigated. The reaction of hexylmagnesium bromide and **6b**, after completion of the ACA and warming to room temperature, afforded the diastereomerically pure *trans*-cyclopropane **8b** in high yield (Table 2, entry 1). Both **7b** and **8b** were obtained with 94% ee. However, precise control of the amount of Grignard reagent is essential to obtain good yields of the cyclopropanes. When less than 1.2 equiv of hexylmagnesium bromide was used, a small but significant

Table 2. Scope of Alkyl Grignard Reagents and Electron-Withdrawing Groups for Tandem ACA–Enolate Trapping toward *trans*-Cyclopropanes^a

entry	R ¹		R ²	product	yield (%)	ee ^b (%)
1	SEt	6b	hexyl	8b	87	94
2	SEt	6b	Me	8g	56	87
3	SEt	6b	Et	8h	67	95
4	SEt	6b	<i>i</i> Pr	8i	89	70
5	SEt	6b	<i>i</i> Bu	8j	91	84
6 ^c	SEt	6b	but-3-enyl	8k	88	94
7 ^d	SEt	6b	(CH ₂) ₃ OrBu	8l	>95 ^d	96
8	SEt	6b	BnCH ₂	8m	92	84
9 ^d	SEt	6b	Ph	8n	50	26
10	C ₁₁ H ₂₃	6e	BnCH ₂	8e	75	96
11	C ₁₁ H ₂₃	6e	Me	8o	87	98
12	OMe	6f	BnCH ₂	8f	68	>95

^a For conditions see Table 1, 1.0 mmol scale. ^b Determined by chiral GC or HPLC. ^c 3.0 mmol scale. ^d Conversion; the cyclopropane proved inseparable by column chromatography from the dimer of the Grignard reagent. ^e 0.5 mmol scale.

amount of acyclic product **7b** was obtained. When more than 1.2 equiv of the Grignard reagent was used, the corresponding cyclopropane bearing a tertiary alcohol was formed, as a result of a subsequent two-fold addition of the Grignard reagent to the thioester moiety.

With the procedure for the synthesis of *trans*-cyclopropanes developed, the scope of the alkyl Grignard reagents for this transformation was explored. The use of both MeMgBr and EtMgBr gave the corresponding volatile *trans*-cyclopropanes **8g** and **8h** with good to excellent ee (entries 2 and 3). The Cu-catalyzed transformation of **6b** with the sterically encumbered Grignard reagents *i*PrMgBr and *i*BuMgBr gave the corresponding cyclopropanes **8i** and **8j** in slightly lower ee but high yield (entries 4 and 5). The catalytic asymmetric synthesis of cyclopropanes using functionalized Grignard reagents includes those substituted with olefin, ether, and aryl groups. These functionalized Grignard reagents gave cyclopropanes **8k**, **8l**, and **8m** with ee's ranging from 84 to 96% (entries 6–8). Finally, when PhMgBr was used the product **8n** was obtained in both low yield and ee (entry 9).

Ketone substrate **6e** gave, upon allowing the reaction mixture to reach room temperature, the corresponding cyclopropane with 96% ee (Table 2, entry 10). The reaction of **6e** with MeMgBr gave cyclopropane **8o** in excellent yield and 98% ee (entry 11). When the reaction mixture of **6f** and phenethylmagnesium bromide was allowed to warm to room temperature, exclusively cyclopropane **8f** was obtained, virtually enantiomerically pure (entry 12).

To illustrate the versatility of the developed method, cyclopropane **8b** was selectively reduced to the aldehyde (see Supporting Information, SI Scheme 1), which is a key intermediate in a reported synthesis¹³ of cascarrillic acid.^{14,15} Furthermore, the reaction of **6b** with heptylmagnesium bromide gave cyclopropane **8p** in 84% yield and 95% ee. Reduction of **8p** to the corresponding aldehyde then gave an intermediate for the synthesis¹⁶ of grenadamide.^{17,18}

In summary, a new and versatile enantioselective synthesis of 4-chloro-3-alkyl-substituted thioesters and ketones and *trans*-1-alkyl-2-substituted cyclopropane esters, thioesters, and ketones has been developed using an extremely simple catalytic system,

based on a commercial chiral ligand, CuI, and Grignard reagents. Application of this methodology in natural product synthesis and other ring structures will be reported in due course.

Acknowledgment. Dedicated to the memory of Dr. M. M. Pollard. We thank M. Bastian for providing substrate **6b** and T. D. Tiemersma-Wegman and M. J. Smith for technical support (GC, HPLC, MS). Financial support from NWO-CW is gratefully acknowledged.

Supporting Information Available: Experimental details, data, and spectra. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) Charette, A. B.; Juteau, H.; Lebel, H.; Molinaro, C. *J. Am. Chem. Soc.* **1998**, *120*, 11943–11952.
- (2) For the asymmetric synthesis of *trans*-1-alkyl-2-substituted cyclopropanes using chiral auxiliaries or substrates, see: (a) Arai, I.; Mori, A.; Yamamoto, H. *J. Am. Chem. Soc.* **1985**, *107*, 8254–8256. (b) Aggarwal, V. K.; Fang, G. Y.; Meek, G. *Org. Lett.* **2003**, *5*, 4417–4420. (c) Risatti, C. A.; Taylor, R. E. *Angew. Chem., Int. Ed.* **2004**, *43*, 6671–6672. (d) Cheeseman, M.; Feuillet, F. J. P.; Johnson, A. L.; Bull, S. D. *Chem. Commun.* **2005**, 2372–2374. (e) Peng, F.; Hall, D. G. *J. Am. Chem. Soc.* **2007**, *129*, 3070–3071. (f) Vega-Pérez, J. M.; Perinán, I.; Iglesias-Guerra, F. *Tetrahedron: Asymmetry* **2009**, *20*, 1065–1072. (g) Cheeseman, M.; Davies, I. R.; Axe, P.; Johnson, A. L.; Bull, S. D. *Org. Biomol. Chem.* **2009**, *7*, 3537–3548. (h) For the formation of *trans*-alkyl substituted cyclopropanes via catalytic asymmetric epoxidation—intramolecular methylene transfer, see: Hardee, D. J.; Lambert, T. H. *J. Am. Chem. Soc.* **2009**, *131*, 7536–7537.
- (3) (a) Fritsch, H.; Leutenegger, U.; Pfaltz, A. *Angew. Chem., Int. Ed. Engl.* **1986**, *25*, 1005–1006. (b) Lo, M. M.-C.; Fu, G. C. *J. Am. Chem. Soc.* **1998**, *120*, 10270–10271. (c) Mazet, C.; Köhler, V.; Pfaltz, A. *Angew. Chem., Int. Ed.* **2005**, *44*, 4888–4891.
- (4) For preparation of these structures via chiral cyclopropanes, see: Lou, Y.; Horikawa, M.; Kloster, R. A.; Hawryluk, N. A.; Corey, E. J. *J. Am. Chem. Soc.* **2004**, *126*, 8916–8918.
- (5) For recent reviews, see: (a) Pellissier, H. *Tetrahedron* **2008**, *64*, 7041–7095. (b) Lebel, H.; Marcoux, J.-F.; Molinaro, C.; Charette, A. B. *Chem. Rev.* **2003**, *103*, 977–1050. (c) Goudreau, S. R.; Charette, A. B. *Angew. Chem., Int. Ed.* **2010**, *49*, 486–488. (d) Doyle, M. P. *Angew. Chem., Int. Ed.* **2009**, *48*, 850–852.
- (6) For recent, selected approaches to substituted cyclopropanes using Au catalysis, see: (a) Watson, I. D. G.; Ritter, S.; Toste, F. D. *J. Am. Chem. Soc.* **2009**, *131*, 2056–2057. Co catalysis, see: (b) Ruppel, J. V.; Gauthier, T. J.; Snyder, N. L.; Perman, J. A.; Zhang, X. P. *Org. Lett.* **2009**, *11*, 2273–2276. Ir catalysis, see: (c) Ichinose, M.; Suematsu, H.; Katsuki, T. *Angew. Chem., Int. Ed.* **2009**, *48*, 3121–3123. B catalysis, see: (d) Zimmer, L. E.; Charette, A. B. *J. Am. Chem. Soc.* **2009**, *131*, 15624–15626. Pd catalysis, see: (e) Liu, W.; Zhu, X.-Z.; Wan, X.-L.; Hou, X.-L. *J. Am. Chem. Soc.* **2009**, *131*, 8734–8735 (synthesis of a *trans*-1-alkyl-2-substituted cyclopropane in 18% yield, 3:1 *trans*:*cis* ratio and 89% ee) Rh catalysis, see: (f) Marcoux, D.; Azzi, S.; Charette, A. B. *J. Am. Chem. Soc.* **2009**, *131*, 6970–6972. (g) DeAngelis, A.; Dmitrenko, O.; Yap, G. P. A.; Fox, J. M. *J. Am. Chem. Soc.* **2009**, *131*, 7230–7231. (h) Lindsay, V. N. G.; Lin, W.; Charette, A. B. *J. Am. Chem. Soc.* **2009**, *131*, 16383–16385. (i) Chuprakov, S.; Kwok, S. W.; Zhang, L.; Lercher, L.; Fokin, V. V. *J. Am. Chem. Soc.* **2009**, *131*, 18034–18035. (j) Grimster, N.; Zhang, L.; Fokin, V. V. *J. Am. Chem. Soc.* **2010**, *132*, 2510–2511. Ru catalysis, see: (k) Xu, Z.-J.; Fang, R.; Zhao, C.; Huang, J.-S.; Li, G.-Y.; Zhu, N.; Che, C.-M. *J. Am. Chem. Soc.* **2009**, *131*, 4405–4417. (l) Ito, J.-I.; Ujije, S.; Nishiyama, H. *Chem.–Eur. J.* **2010**, *16*, 4986–4990.
- (7) For the non-asymmetric addition of Grignard reagents, see: (a) Ratney, R. S.; English, J., Jr. *J. Org. Chem.* **1960**, *25*, 2213–2215. (b) Torii, S.; Tanaka, H.; Nagai, Y. *Bull. Chem. Soc. Jpn.* **1977**, *50*, 2825–2826. (c) Eisch, J. J.; Galle, J. E. *J. Org. Chem.* **1979**, *44*, 3277–3279. Li reagents, see: (d) Cooke, M. P., Jr.; Jaw, J. Y. *J. Org. Chem.* **1986**, *51*, 758–760.
- (8) These compounds have been observed as a side product: Lee, Y.; Hoveyda, A. H. *J. Am. Chem. Soc.* **2006**, *128*, 15604–15605.
- (9) (a) Lu, Z.; Ma, S. *Angew. Chem., Int. Ed.* **2008**, *47*, 258–297. (b) Harutyunyan, S. R.; den Hartog, T.; Geurts, K.; Minnaard, A. J.; Feringa, B. L. *Chem. Rev.* **2008**, *108*, 2824–2852. (c) Alexakis, A.; Bäckvall, J. E.; Krause, N.; Pàmies, O.; Diéguez, M. *Chem. Rev.* **2008**, *108*, 2796–2823. (d) den Hartog, T.; Maciá, B.; Minnaard, A. J.; Feringa, B. L. *Adv. Synth. Catal.* **2010**, *352*, 999–1013.
- (10) (a) Feringa, B. L.; Badorrey, R.; Peña, D.; Harutyunyan, S. R.; Minnaard, A. J. *Proc. Natl. Acad. Sci. U.S.A.* **2004**, *101*, 5834–5838. (b) López, F.; Harutyunyan, S. R.; Minnaard, A. J.; Feringa, B. L. *J. Am. Chem. Soc.* **2004**, *126*, 12784–12785. (c) López, F.; Harutyunyan, S. R.; Minnaard, A. J.; Feringa, B. L. *Angew. Chem., Int. Ed.* **2005**, *44*, 2752–2756. (d) DesMazery, R.; Pullez, M.; López, F.; Harutyunyan, S. R.; Minnaard, A. J.; Feringa, B. L. *J. Am. Chem. Soc.* **2005**, *127*, 9966–9967. (e) Wang, S.-Y.; Ji, S.-J.; Loh, T.-P. *J. Am. Chem. Soc.* **2007**, *129*, 276–277. (f) Maciá, B.; Geurts, K.; Fernández-Ibáñez, M. A.; ter Horst, B.; Minnaard, A. J.; Feringa, B. L. *Org. Lett.* **2007**, *9*, 5123–5126. (g) Wang, S.-Y.; Lum, T.-K.; Ji, S.-J.; Loh, T.-P. *Adv. Synth. Catal.* **2008**, *350*, 673–677. (h) Jerphagnon,

- T.; Pizzuti, M. G.; Minnaard, A. J.; Feringa, B. L. *Chem. Soc. Rev.* **2009**, 38, 1039–1075.
- (12) The low yield of **8a** is mainly due to incomplete conversion of **6a** and the competitive formation of ethyl thiocrotonate via halogen–magnesium exchange and subsequent protonation upon workup.
- (13) Cheeseman, M.; Bull, S. D. *Synlett* **2006**, 1119–1121.
- (14) Roberts, I. O.; Baird, M. S.; Liu, Y. *Tetrahedron Lett.* **2004**, 45, 8685–8686.
- (15) Isolation: (a) Thoms, H. *Verh. Vers. Deutsch. Naturf. Aerzte* **1899**, 648–652. (b) Thoms, H.; Fendler, G. *Arch. Pharm.* **1900**, 238, 671–690. Structure elucidation: (c) Motl, O.; Amin, M.; Sedmera, P. *Phytochemistry* **1972**, 11, 407–408.
- (16) Al Dulayymi, J. R.; Baird, M. S.; Jones, K. *Tetrahedron* **2004**, 60, 341–345.
- (17) (a) Green, R.; Cheeseman, M.; Duffill, S.; Merritt, A.; Bull, S. D. *Tetrahedron Lett.* **2005**, 46, 7931–7934. (b) Avery, T. D.; Culbert, J. A.; Taylor, D. K. *Org. Biomol. Chem.* **2006**, 4, 323–330.
- (18) Isolation: Sitachitta, N.; Gerwick, W. H. *J. Nat. Prod.* **1998**, 61, 681–684.

JA105704M